

# Your Body Already Knows

Sean Morin | 2748684 Alberta Ltd. | April 7, 2026

---

## 1. The 23-Minute Cost of Decoupling

---

The average time to resume focused work after a single interruption is 23 minutes and 15 seconds. Workers sustain a task for approximately 3 minutes before switching, maintain a working sphere for 12 minutes, and 57% of tasks are interrupted before completion [1].

Creative flow requires transient hypofrontality — decreased activity in the superior frontal gyri, the brain's executive control regions. EEG recordings of jazz musicians during improvisation confirmed that high-flow states correlate with reduced superior frontal activity and increased left-hemisphere auditory and sensorimotor processing [2].

Removing background audio produces the same resumption penalty as a visual interruption. Silence is not neutral. It is the removal of a synchronization signal.

**Therefore:** The interruption cost is not caused by distraction. It is caused by the brain's oscillatory rhythms losing synchronization with the work environment. Recovery takes 23 minutes because the rhythms must recouple from scratch.

---

## 2. The 100-120 BPM Pacemaker

---

Rhythmic auditory stimulation at 100-120 BPM activates the basal ganglia independently of conscious intent — the tempo range at which humans universally experience involuntary motor entrainment to a beat.

The Optimal Stimulation Theory of ADHD proposes that hyperactive and inattentive behaviors reflect attempts to compensate for chronic cortical

underarousal. External rhythmic input provides the stimulation the dopaminergic system cannot generate internally, improving inhibitory control and task initiation [3].

Music intervention in chronic pain patients significantly improves heart rate variability (HRV) — a core biomarker of autonomic nervous system stability — while simultaneously reducing self-reported pain intensity. A randomized controlled study of 37 chronic pain patients demonstrated measurable HRV improvement and prefrontal hemodynamic changes after seven days of music intervention [4].

When heart rate and gait rhythm achieve 1:1 phase-lock — a state called cardiac-locomotor synchronization — stroke volume increases, blood pressure variability decreases, and oxygen uptake drops [5].

**Therefore:** 100-120 BPM sits near the resonant frequency of the human cardiac-locomotor coupling system. It simultaneously functions as a dopaminergic pacemaker for ADHD and an autonomic stabilizer for chronic pain. This is not a music preference. It is a biological frequency match.

---

### 3. The 45% Overlap

---

In a study of 123 fibromyalgia patients screened using the WHO Adult ADHD Self-Report Scale, 44.7% met diagnostic criteria for ADHD [6].

A systematic review of 71 studies confirmed significant changes in both sympathetic and parasympathetic autonomic nervous system function during pain, independent of induction method [7].

Both conditions involve dysregulation of the same neurotransmitters — dopamine and norepinephrine. ADHD is characterized by failure to synchronize cortical rhythms with environmental temporal structure. Fibromyalgia is characterized by autonomic rhythm instability — cardiac and locomotor rhythms fail to maintain synchronization, producing HRV disruption and central pain amplification.

**Therefore:** These are not two separate diseases with coincidental comorbidity. They are the same coupling failure in different subsystems — cortical-environmental in ADHD, autonomic-locomotor in fibromyalgia — sharing the same neurotransmitter infrastructure.

This explains why stimulant medication sometimes reduces chronic pain. Why exercise improves ADHD symptoms. Why rhythm — drumming, dancing, music — helps both. Each provides an external oscillation the desynchronized system can lock onto.

---

## 4. Why Tracking Apps Fail

---

Patient symptom diaries achieve less than 50% accuracy across chronic conditions. Patient intuition about pre-flare states remains high despite diary failure. The knowledge exists. The reporting mechanism is broken.

Pre-flare awareness is embodied — it manifests as changes in movement patterns, heart rate variability, and gait cadence. It is not verbal. As Polanyi established: "We know more than we can tell" [10].

Directed attention is a depletable resource. Restorative environments — those engaging involuntary "soft fascination" rather than directed attention — measurably restore cognitive capacity [9]. Interfaces that demand conscious cognitive effort drain the same energy pool that chronic illness already depletes.

Spoon Theory describes the finite energy budget of chronic illness [11]. Any interface requiring cognitive load to operate consumes the resource it claims to help manage.

**Therefore:** Symptom tracking apps fail not because patients are non-compliant, but because the capture mechanism is incompatible with the data format. The body's signal is rhythmic and embodied. The form is static and verbal. The translation destroys the signal.

---

## 5. The Auditory Advantage

---

Auditory simple reaction time averages 284 milliseconds. Visual simple reaction time averages 331 milliseconds [8]. The auditory pathway from inner ear to temporal cortex is anatomically shorter and requires fewer synaptic relays than the visual pathway from retina to occipital cortex.

Auditory biofeedback produces superior long-term retention compared to visual biofeedback. Visual feedback creates dependency. Auditory feedback allows internalization through proprioception.

**Therefore:** Sound is the fastest sensory pathway to the motor system. It is the only modality fast enough to maintain continuous synchronization with biological rhythms without requiring conscious attention.

---

## 6. What This Means

---

The research cited in this brief comes from auditory neuroscience, cardiology, rehabilitation medicine, chronic pain management, ADHD psychiatry, phenomenology, and human-computer interaction. These fields do not read each other's journals.

When read together, they describe one phenomenon: the human body is a system of coupled oscillators. Focus, health, and creative flow are states of synchronization. Distraction, pain amplification, and executive dysfunction are states of desynchronization.

Music is not background noise. It is an external pacemaker. Silence is not neutral. It is the removal of a coupling signal. Chronic pain and ADHD are not separate diseases. They are the same coupling failure in different subsystems.

---

## References

---

1. Mark, G., Gonzalez, V.M., & Harris, J. (2005). No task left behind? Examining the nature of fragmented work. *Proceedings of the SIGCHI Conference on Human Factors in Computing Systems (CHI '05)*, 321-330. ACM. <https://doi.org/10.1145/1054972.1055017>
2. Rosen, D., Oh, Y., Chesebrough, C., Zhang, F.Z., & Kounios, J. (2024). Creative flow as optimized processing: Evidence from brain oscillations during jazz improvisations by expert and non-expert musicians. *Neuropsychologia*, 196, 108824. <https://doi.org/10.1016/j.neuropsychologia.2024.108824>
3. Zentall, S.S. & Zentall, T.R. (1983). Optimal stimulation: A model of disordered activity and performance in normal and deviant children. *Psychological Bulletin*, 94, 446-471. See also: Zentall, S.S. (1975). Optimal stimulation as theoretical basis of hyperactivity. *American Journal of Orthopsychiatry*, 45(4), 549-563. <https://doi.org/10.1111/j.1939-0025.1975.tb01185.x>
4. Chen, Z., Wang, Y., Wang, R., Chai, Y., Zhang, Y., Deng, Y., & Pan, J. (2022). Effect of music intervention on subjective scores, heart rate variability, and prefrontal hemodynamics in patients with chronic pain. *Frontiers in Human Neuroscience*, 16, 1057290. <https://doi.org/10.3389/fnhum.2022.1057290>
5. Niizeki, K., Kawahara, K., & Miyamoto, Y. (1993). Interaction among cardiac, respiratory, and locomotor rhythms during cardiocomotor synchronization. *Journal of Applied Physiology*, 75(4), 1815-1821. <https://doi.org/10.1152/jappl.1993.75.4.1815>
6. van Rensburg, R., Meyer, H.P., Hitchcock, S.A., & Schuler, C.E. (2018). Screening for adult ADHD in patients with fibromyalgia syndrome. *Pain Medicine*, 19(9), 1825-1831. <https://doi.org/10.1093/pm/pnx275>

7. Forte, G., Troisi, G., Pazzaglia, M., De Pascalis, V., & Casagrande, M. (2022). Heart rate variability and pain: A systematic review. *Brain Sciences*, 12(2), 153. <https://doi.org/10.3390/brainsci12020153>
8. Shelton, J. & Kumar, G.P. (2010). Comparison between auditory and visual simple reaction times. *Neuroscience & Medicine*, 1(1), 30-32. <https://doi.org/10.4236/NM.2010.11004>
9. Kaplan, S. (1995). The restorative benefits of nature: Toward an integrative framework. *Journal of Environmental Psychology*, 15(3), 169-182. [https://doi.org/10.1016/0272-4944\(95\)90001-2](https://doi.org/10.1016/0272-4944(95)90001-2)
10. Polanyi, M. (1966). *The Tacit Dimension*. University of Chicago Press.
11. Miserandino, C. (2003). The spoon theory. [butyoudontlooksick.com](http://butyoudontlooksick.com).